

# Taxonomic Diversity of Fungi Associated with Some PCN Populations from Serbia

Violeta Oro<sup>1</sup>, Jelena Bošković<sup>2</sup>, Slobodan Milenković<sup>2</sup> and Solveig Tosi<sup>3</sup>

<sup>1</sup>*Institute for Plant Protection and Environment, Department of Plant Pathology, Nematology Laboratory, Teodora Drajzera 9, 11000 Belgrade, Serbia (viooro@yahoo.com)*

<sup>2</sup>*Megatrend University, Faculty of Biofarming, Maršala Tita 39, 24300 Bačka Topola, Serbia*

<sup>3</sup>*University of Pavia, Department of Earth Science and Environment, Mycology Laboratory, Pavia, Italy*

*Received: October 10, 2011*

*Accepted: February 24, 2012*

## SUMMARY

Increased content of pesticides in food chain resulted in using microorganisms as agents of biological control. The potato cyst nematodes (PCN) - *Globodera pallida* and *G. rostochiensis* belong to the group of the most important parasites - the quarantine organisms. The external and internal area of a cyst harbor numerous fungal and bacterial species. The aim of this study was to identify antagonistic fungi associated with some PCN populations from Serbia. Fungal antagonists of potato cyst nematodes have not been previously investigated in our country. The diversity of PCN fungal antagonists is not reflected only at the species level but also at the level of higher taxonomic categories.

**Keywords:** Potato Cyst Nematodes (PCN); Fungi; Diversity; Biological control; Serbia

## INTRODUCTION

Increased public concern over the significant presence of pesticides in food chain and its consequences resulted in the development of biological agents as alternative methods of control invasive organisms.

The potato cyst nematodes (PCN) - *Globodera pallida* and *G. rostochiensis* belong to the group of the most important parasites-the quarantine organisms and their control is complex due to their long persistence in soil.

The fungal antagonists of nematodes consist of a great variety of organisms which include the nematode-trapping or predacious fungi, endoparasitic

fungi, parasites of nematode eggs and cysts and fungi which produce metabolites toxic to nematodes. The fungal spore has undergone some remarkable adaptation and specialization to capture or penetrate nematodes. Among the lower fungi, motile zoospores appear to have positive tropism toward nematodes. *Catenaria anguillulae* spores most often accumulate around the natural body orifices of the nematode cuticle (Mankau, 1980). *Nematophthora gynophila* causes total destruction of *Heterodera avenae* cysts in less than seven days (Kerry, 1980). *Verticillium chlamyosporium*, *Acremonium strictum* and *Fusarium oxysporum* are the main parasites in eggs of *H. schachtii* (Dackman and Nordbring-Hertz, 1985).

Flavipin, a low molecular weight compound, a metabolite of the fungus *Chaetomium globosum* is responsible for most of the nematode-antagonistic activity (Nita et al., 2002). Combined application of *Paecilomyces lilacinus* and *Monacrosporium lysipagum* reduced 65% of *Heterodera avenae* cysts (Khan et al., 2006). There is a commercialised biopesticide based on *Paecilomyces lilacinus* (Kerry, 2000). In our country, fungal antagonists of potato cyst nematodes have not been investigated so far. The aim of the study was to identify taxonomic diversity of fungi associated with some PCN populations from Serbia.

## MATERIAL AND METHODS

Mycobiota of *Globodera* cysts from Kladnica, Šanac, Gojna Gora and Milatovići populations which are morphologically and molecularly identified (Oro et al., 2010) was investigated. The cysts (3) and eggs of a single cyst were placed on potato-dextrose agar (PDA), containing antibiotics (bensylpenicillin K + bensylpenicillin-procaine: 200.000 i.u.+600.000 i.u., 200 mg/l), and incubated for two weeks at 25 a°C. The cysts and eggs were previously disinfected according to Heungens et al., 1996.

For identification purpose fungi were maintained on MEA (Malt Extract Agar) at room temperature. The isolates were identified by morpho-dimensional analysis using slide-cultures of the fungi. Only *Mycelia Sterilia* was identified by biomolecular analysis.

## RESULTS AND DISCUSSION

The external and internal area of a cyst harbor numerous fungal and bacterial species competing for substrate. In the process of identification, the following fungal species were found: *Fusarium oxysporum* on cysts of the Kladnica population, *Oxyporus latemarginatus* on eggs of the same population, *Botrytis cinerea* and *Aspergillus fumigatus*, on eggs of the Šanac population, *Geotrichum candidum* on cysts of the Gojna Gora population, *Oxyporus latemarginatus* on eggs of the same population, *Geotrichum candidum* on cysts of the Milatovići population and *Aspergillus fumigatus* was isolated on eggs of the Milatovići population. The found species are harmless saprobes and they are relatively abundant members of the soil microbial community. Nearly all of them performed antagonistic activity toward potato cyst nematodes and

have the potential to be used as biocontrol agents (results are not shown). The morphological view of isolated fungi is presented in photos.

### *Fusarium oxysporum* Schlecht. Fr.

Fungi

Dikarya

Ascomycota

Pezizomycotina

Sordariomycetes

Hypocreales

Nectriaceae

*Fusarium* Teleomorph: *Gibberella*

*Fusarium* is a large genus of filamentous fungi widely distributed in soil and in association with plants (Ohara et al., 2004).



Figure 1. Mycelium of *Fusarium oxysporum* (PDA)



Figure 2. *Fusarium oxysporum* (LM 400X)



Figure 3. Mycelium of *Oxyporus latemarginatus* (PDA)



Figure 4. *Oxyporus latemarginatus* (Di 13X)

Many *F. oxysporum* isolates appear to be host specific, which has resulted in the subdivision of the species into *formae speciales* and races that reflect the apparent plant pathogenic specialization. *Fusarium oxysporum* and its various *formae speciales* have been characterized as causing the following symptoms: vascular wilt, yellows, corm rot, root rot, damping-off etc. (Kim et al., 2002). In solid media culture, such as potato dextrose agar (PDA), different special forms of *F. oxysporum* can have varying appearances. On PDA, growth is rapid and the white aerial mycelium may become tinged with purple or be submerged by the blue color of the sclerotia or by the cream to tan to orange sporodochia when they are abundant (Nelson et al., 1983; Leslie and Summerell, 2006). *Fusarium oxysporum* produces three types of asexual spores: microconidia, macroconidia, and chlamydospores (Ruiz-Roldan et al., 2010). Mycelium of *Fusarium oxysporum* on PDA and under a light microscope (LM) are shown in figures 1 and 2 respectively.

### *Oxyporus latemarginatus* (Durieu and Mont.) Donk

Fungi  
Dikarya  
Basidiomycota  
Agaricomycotina  
Agaricomycetes  
Polyporales  
Polyporaceae  
*Oxyporus*

*Oxyporus latemarginatus* is widely distributed in Europe; it induces stem and root rot of *Fagus* and *Quercus* species. The fungus colonizes and survives on dead wood or roots in the soil, forming white mycelial mats and dirty white to pale yellow fruiting bodies, annual, resupinate, rather soft when fresh, 2-5 (10) mm thick (Ryvard and Gilbertson, 1994). Mycelium of *Oxyporus latemarginatus* on PDA and under a dissecting microscope (Di) are shown in figures 3 and 4 respectively.

### *Botrytis cinerea* (De Bary) Whetzel

Fungi  
Dikarya  
Ascomycota  
Pezizomycotina  
Leotiomycetes  
Helotiales  
Sclerotiniaceae  
*Botrytis* Teleomorph: *Botryotinia*

*Botrytis cinerea* has a worldwide distribution but occurs mainly in humid temperate and subtropical regions; it is very common in the phylloplane of plants but does not attack healthy leaves; it can be facultative parasite on a wide range of plants causing blight or rot of leaves, flowers and fruits, the so-called gray mold (Domsch et al., 1980).



Figure 5. Mycelium of *Botrytis cinerea* (PDA)



Figure 6. *Botrytis cinerea* (LM 400X)




Figure 7. Mycelium of *Aspergillus fumigatus* (PDA)



Figure 8. *Aspergillus fumigatus* (LM 400X)

*Botrytis cinerea* causes infections characterized by rapid destruction (maceration, necrosis) of the tissues of its plant host as it colonizes it (necrotrophy).

The fungus overwinters as mycelia or black resting structures called sclerotia that may be evident on dead plant tissue in late summer. Sclerotia germinate

in the spring, or mycelium grows out of infected debris and conidia (infectious spores) develop (Williamson et al., 2007).

Mycelium of *Botrytis cinerea* on PDA and under a light microscope (LM) are shown in figures 5 and 6 respectively.

### ***Aspergillus fumigatus* Fresenius**

Fungi

Dikarya

Ascomycota

Pezizomycotina

Eurotiomycetes

Eurotiales

Trichocomaceae

*Aspergillus* Teleomorph: *Emericella*

*Aspergillus fumigatus*, a saprotroph widespread in nature, is typically found in soil and decaying organic matter, where it plays an essential role in carbon and nitrogen cycling. Colonies of the fungus produce from conidiophores thousands of minute grey-green conidia (2-3  $\mu\text{m}$ ) that readily become airborne. For many years, *A. fumigatus* was thought to reproduce only asexually, as neither mating nor meiosis had ever been observed. O’Gorman et al., 2009 proved *A. fumigatus* possess a fully functional sexual reproductive cycle. The fungus is capable of growth at 37° C up to 50° C, with conidia surviving at 70° C. Mycelium of *Aspergillus fumigatus* on PDA and under a light microscope (LM) are shown in figures 7 and 8 respectively.

### ***Geotrichum candidum* Link**

Fungi

Dikarya

Ascomycota

Saccharomycotina

Saccharomycetes

Saccharomycetales

Endomycetaceae

*Geotrichum* Teleomorph: *Dipodascus*

*Geotrichum candidum* is an extremely common fungus with a worldwide distribution, occurring mainly in soil, air, water, sewage, various plant substrates and very frequently on milk and milk products (Domsch et al., 1980). It can be a plant pathogen that causes sour rot on peach, nectarine, tomato and carrot. Species of the genus *Geotrichum* produce chains of hyaline, smooth, one-celled, subglobose to cylindrical, slimy arthroconidia

(Gente et al., 2006). Mycelium of *Geotrichum candidum* on PDA and under a light microscope (LM) are shown in figures 9 and 10 respectively.



Figure 9. Mycelium of *Geotrichum candidum* (PDA)




Figure 10. *Geotrichum candidum* (LM 400X)

The mode of antagonistic action of fungi on potato cyst nematodes is very complex and involves numerous toxins, metabolites, lytic enzymes and other anti-helminthic compounds.

The main toxins produced by *Fusarium* species are fumonisins and trichothecenes (Embaby and Abdel-Galil, 2006). *Fusarium oxysporum* is a well known nematode antagonist used in many surveys (Morgan-Jones and Kabana, 1981; Bertrand et al., 2000; Athman et al., 2006; Daami-Remadi et al., 2009). Some strains are reported to be mycoparasites (Gupta et al., 1979), or parasites of weeds (Amsellem et al., 2001) and insects (Warma and Tandam, 1996). The fungus is a member of Nectriaceae family of Ascomycota phylum.

*Oxyporus latemarginatus* produces volatile, anti-fungal compound 5-pentyle-2-furaldehyde which is effective

against *Alternaria alternata*, *Botrytis cinerea*, *Colletotrichum gloeosporoides*, *Magnaporthe grisea*, *Rhizoctonia solani*, *Fusarium oxysporum* f.sp. *lycopersici* (Lee et al., 2008). Pentyle-furaldehyde (PTF) was first reported as a nematocidal metabolite against *Aphelenchoides besseyi* from *Irpex lacteus* IFO5367 (Hayashi et al. 1981). Mayer et al. (1996) reported that PTF produced by an Ascomycete exhibits nematocidal activity against *Caenorhabditis elegans* and *Meloidogyne incognita*. *Oxyporus* belongs to Polyporaceae family and it is the only representative of Basidiomycota.

*Botrytis cinerea* utilizes a large range of pathogenic factors (lytic enzymes, activated oxygen forms, toxins, plant hormones) to attack its host plants but it was not reported as nematode antagonist. The fungus is a member of Sclerotiniaceae family of Ascomycota.

When the fermentation broth of *Aspergillus fumigatus* was screened, a number of indolic alkaloids with antimetabolic properties were discovered such as tryprostatins, with spirotryprostatin B being of special interest as an anticancer drug (Cui et al., 1996). *Aspergillus fumigatus* grown on certain building materials can produce genotoxic and cytotoxic mycotoxins, such as gliotoxin (Nieminen et al., 2002), it can also produce antibiotics such as sideramines fusigen, ferricrocin and fumigacin (Domsch et al., 1980). *Aspergillus fumigatus* had immobilized 51-75% juvenils of *Meloidogyne* spp. in three days (Mubyana-John and Wright, 2011). *Aspergillus* belongs to Trichocomaceae family of Ascomycota.

Insecticidal neofrapeptins were isolated from *Geotrichum candidum* (Nagaraj et al., 2001). High egg parasitization (72%) of *Meloidogyne incognita* was observed by *Geotrichum candidum* (Goswami and Mittal, 2002). The fungus belongs to Endomycetaceae family or sometimes it is classified in Saccharomycetaceae, family of yeasts, within Ascomycota.

The process of cyst colonisation by fungi is enhanced by cystogenesis.

In shape, cysts are spherical sacs in which 500-600 eggs can be laid. There are specific ecological conditions within the cyst (temperature, relative humidity, pH, presence of different chemical compounds) which make it an ideal medium for microbial growth. Generally, eggs and larvae are not infected while cysts are intact. In process of hatching, the first larva appears leaving the vulva open and allowing fungal colonization. When detached from root tissue, mature cysts often lack a head leaving the cyst open on the other side, so that in this way antagonistic microorganisms can invade from both directions. In this study, different fungal species colonize external and internal side of a cyst. Although the majority of mentioned

fungal species are occasionally found in association with other nematode species, *Oxyporus latemarginatus* is for the first time found to be associated with PCNs. The diversity of PCN fungal antagonists is not reflected only at the species level but also at the level of higher taxonomic categories i.e. in existence of different phyla such as Ascomycota and Basidiomycota. Thus, the cyst can be considered as specific „microcosm“ containing a spectrum of different fungi inside and outside of the cyst.

## ACKNOWLEDGMENT

This study was supported by the Ministry of Education and Science Grants TR 31018 and I I I 46007.

## REFERENCES

- Amsellem, Z., Kleifeld, Y., Kerenyi, Z., Hornok, L., Goldvasser, Y. and Gressel, J.:** Isolation, identification, and activity of mycoherbicidal pathogens from juvenile broomrape plants. *Biological Control*, 21: 274-284, 2001.
- Athman, S.Y., Dubois, T., Coyne, D., Gold, C.S., Labuschagne, N. and Viljoen, A.:** Effect of endophytic *Fusarium oxysporum* on host preference of *Radopholus similis* to tissue culture banana plants. *Journal of Nematology*, 38(4): 455-460, 2006.
- Bertrand, B., Nuñez, C. and Sarah, J.L.:** Disease complex in coffee involving *Meloidogyne arabicida* and *Fusarium oxysporum*. *Plant Pathology*, 49: 383-388, 2000.
- Cui, C-B., Kakeya, H. and Osada, H.:** Novel mammalian cell cycle inhibitors, spirotryprostatin A and B, produced by *Aspergillus fumigatus*, which inhibit mammalian cell cycle at G2/M phase. *Tetrahedron*, 52(39): 12651-12666, 1996.
- Dackman, C. and Nordbring-Hertz, B.:** Fungal parasites of the cereal cyst nematode *Heterodera avenae* in southern Sweden. *Journal of Nematology*, 17(1): 50-55, 1985.
- Daami-Remadi, M., Sayes, S., Horigue-Raouani, N. and Hlaoua-Ben Hassine, W.:** Effects of *Verticillium dahliae* Kleb., *Fusarium oxysporum* Schlecht. f. sp. *tuberosi* Snyder, Hansen and *Meloidogyne javanica* (Treb.) Chitwood inoculated individually or in combination on potato growth, wilt severity and nematode development. *African Journal of Microbiology Research*, 3(10): 595-604, 2009.
- Domsch, K.H., Gams, W. and Anderson, T.-H.:** *Compendium of Soil Fungi*. Vol. 1. Academic Press, Great Britan, 1980, pp. 150-155.
- Eapen, S.J., Beena, B. and Ramana, K.V.:** Tropical soil microflora of spice-based cropping systems as potential antagonists of root-knot nematodes. *Journal of Invertebrate Pathology*, 88: 218-225, 2005.
- Embaby, E.M. and Abdel-Galil, M.M.:** Seed borne fungi and mycotoxins associated with some legume seeds in Egypt. *Journal of Applied Sciences Research*, 2(11): 1064-1071, 2006.
- Gente, S., Sobier, D., Coton, E., Dubamel, C. and Gueguen, M.:** Identification of *Geotrichum candidum* at the species and strain level: proposal for a standardized protocol. *Journal of Industrial Microbiology and Biotechnology*, 33: 1019-1031, 2006.
- Goswami, B.K. and Mittal, A.:** Effect of some fungal bioagents on root-knot nematode *Meloidogyne incognita* infecting brinjal (Abstract). *Pakistan Journal of Nematology*, 20(1): 55-59, 2002.
- Gupta, R.C., Upadhyay, R.S. and Rai, B.:** Hyphal parasitism and chlamydospore formation by *Fusarium oxysporum* on *Rhizoctonia solani*. *Mycopathologia*, 67: 147-152, 1979.
- Hayashi, M., Wada, K. and Munakata, K.:** New nematocidal metabolites from a fungus, *Irpex lacteus*. *Agricultural and Biological Chemistry*, 45: 1527-1529, 1981.
- Heungens, K., Mugniery, D., Van Montag, M., Gheysen, G. and Niebel, A.:** A method to obtain disinfected *Globodera* infective juveniles directly from cysts. *Fundamental and Applied Nematology*, 19(1): 91-93, 1996.
- Kerry, B.R.:** Biocontrol: fungal parasites of female cyst nematodes. *Journal of Nematology*, 12: 253-259, 1980.
- Kerry, B.R.:** Rhizosphere interactions and the exploitation of microbial agents for the biological control of plant-parasitic nematodes. *Annual Review of Phytopathology*, 38: 423-441, 2000.
- Khan, A., Williams, K.L. and Nevalainen, H.K.M.:** Infection of plant-parasitic nematodes by *Pacelomyces lilacinus* and *Monacrosporium lysipagum*. *BioControl*, 51: 659-687, 2006.
- Kim, W.G., Byung, D.L., Kim, W.S. and Cho, W.D.:** Root rot of moth orchid caused by *Fusarium* spp. *Plant Pathology Journal*, 18(4): 225-227, 2002.
- Lee, S.O., Kim, H.Y., Choi, G.J., Lee, H.B., Jang, K.S., Choi, Y.H. and Kim, J.-C.:** Mycofumigation with *Oxyporus latemarginatus* EF069 for control of postharvest apple decay and *Rhizoctonia* root rot on moth orchid. *Journal of Applied Microbiology*, 106: 1213-1219, 2009.
- Leslie, J.F. and Summerell, B.A.:** *Fusarium* Laboratory Manual. Blackwell Publishing, Ames, Iowa, USA, 2006, pp. 388.
- Mankau, R.:** Biocontrol: Fungi as nematode control agents. *Journal of Nematology*, 12(4): 244-252, 1980.
- Mayer, A., Köpke, B., Anke, H. and Sterner, O.:** Dermatolactone, a cytotoxic fungal sesquiterpene with a novel skeleton. *Phytochemistry*, 43(2): 375-376, 1996.

- Morgan-Jones, G. and Rodriguez-Kabana, R.:** Fungi associated with cysts of *Heterodera glycines*. *Nematropica*, 11(1): 69-74, 1981.
- Mubyana-John, T. and Wright, M.G.:** Fungi and their use in the possible control of nematodes in Botswana soils. *African Journal of Plant Science and Biotechnology*, 5: 33-40, 2011.
- Nagaraj, G., Uma, M.V., Shivayogi, M.S. and Balaram, H.:** Antimalarial activities of peptide antibiotics isolated from fungi. *Antimicrobial Agents and Chemotherapy*, 45: 145-149, 2001.
- Nelson, P.E., Toussoun, T.A. and Marasas, W.F.O.:** *Fusarium* Species. An illustrated Manual for Identification. The Pennsylvania State University Press, USA, 1983, pp. 193.
- Nieminen, S.M., Mäki-Paakkanen, J., Hirvonen, M-R., Roponen, M. and Von Wright, A.:** Genotoxicity of gliotoxin, a secondary metabolite of *Aspergillus fumigatus*, in a battery of short-term test systems. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 520(1-2): 161-170, 2002.
- Nitao, J.K., Meyer, S.L.F., Oliver, J.E., Schmidt, W.F. and Chitwood D.J.:** Isolation of flavipin, a fungus compound antagonistic to plant-parasitic nematodes. *Nematology*, 4(1): 55-63, 2002.
- O’Gorman, C.M., Fuller, H.T. and Dyer, P.S.:** Discovery of a sexual cycle in the opportunistic fungal pathogen *Aspergillus fumigatus*. *Nature*, 457: 471-475, 2009.
- Obara, T., Inoue, I., Namiki, F., Kunob, H. and Tsuge, T.:** *REN1* is required for development of microconidia and macroconidia, but not of chlamydospores, in the plant pathogenic fungus *Fusarium oxysporum*. *Genetics*, 166: 113-124, 2004.
- Oro, V., Ivanović, Ž., Nikolić, B., Barszi, L., Radivojević, M. and Jovčić, B.:** Morphological and molecular identification of potato cyst nematode populations in Serbia. *Archives of Biological Sciences*, 62(3): 747-754, 2010.
- Ruiz-Roldan, M.C., Köbli, M., Roncero, M.I.G., Philippesen, P., Di Pietro, A. and Espeso, E.A.:** Nuclear dynamics during germination, conidiation, and hyphal fusion of *Fusarium oxysporum*. *Eukaryotic Cell*, 9(8): 1216-1224, 2010.
- Ryvard, L. and Gilbertson, R.L.:** European Polypores. Part 2. Synopsis Fungorum 7, Fungiflora, Oslo, Norway, 1994, pp. 442-443.
- Warma, A. and Tandam, B.K.:** Pathogenicity of three entomogenous fungi against insect pests of sugarcane. *Journal of Biological Control*, 10: 87-91, 1996.
- White, T.J., Bruns, T., Lee, S. and Taylor, J.W.:** Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: *PCR Protocols: a Guide to Methods and Applications* (Innis M.A., Gelfand D.H., Sninsky J.J., White T.J., eds). Academic Press, San Diego, USA, 1990, pp. 315-322.
- Williamson, B., Tudzinsky, B., Tudzinsky, P. and Van Kan, J.A.L.:** *Botrytis cinerea*: the cause of grey mold disease. *Molecular Plant Pathology*, 8(5): 561-580, 2007.

## Taksonomski diverzitet gljivica povezanih sa nekim populacijama CNK iz Srbije

### REZIME

Povećan sadržaj pesticida u lancu ishrane je rezultirao korišćenjem mikroorganizama kao agenasa biološke kontrole. Cistolike nematode krompira (CNK) – *Globodera pallida* i *G. rostochiensis* spadaju u grupu najznačajnijih parazita – karantinskih organizama. Spoljašnjost i unutrašnjost ciste krije brojne vrste gljiva i bakterija. Cilj ovog rada je bio da se identifikuju gljivice u asocijaciji sa nekim populacijama CNK iz Srbije. U našoj zemlji fungalni antagonisti cistolikih nematoda krompira nisu bili istraživani do sada. Diverzitet fungalnih antagonista CNK se ne reflektuje samo na specijskom nivou, već i na nivou viših taksonomskih kategorija.

**Ključne reči:** Cistolike nematode krompira (CNK); gljivice; diverzitet; biološka kontrola; Srbija